arXiv:hep-th/9806145v1 17 Jun 1998

Hidden conformal symmetry of a massive scalar field in AdS$_2$ [*]

J. Cruz [†]
Center for Theoretical Physics
Laboratory for Nuclear Science
and Department of Physics
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139
and
Departamento de Fisica Teorica and IFIC
Centro Mixto Universidad de Valencia-CSIC
Facultad de Fisica, Universidad de Valencia
Burjassot-46100, Valencia, Spain

(MIT-CTP-2752, hep-th/9806145. June 1998)

Abstract

We show that a massive scalar field on a 2-dimensional Anti-De Sitter space (AdS$_2$) can be mapped by means of a time-dependent canonical transformation into a massless free field theory when the mass square of the field is equal to minus the curvature of the background metric. We also provide the (hidden) conformal symmetry of the massive scalar field.

PACS number(s): 04.60.Kz, 04.60.Ds
Keywords: Anti-De Sitter space, Canonical Transformations, Conformal Symmetry.

Submitted to: *Phys. Lett. B*

[†] Email addresses: `cruz@ctpa03.mit.edu`
`cruz@lie.uv.es`

[*] This work is supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under cooperative research agreement #DF-FC02-94ER40818.

The Liouville action

$$S_L = -\frac{1}{2}\int d^2x \left[\partial_\mu\phi\partial^\mu\phi + \frac{\lambda^2}{\beta^2}e^{\beta\phi}\right] , \tag{1}$$

defined over the 2-dimensional Minkowski space, is not invariant under conformal transformations when the field ϕ transforms in the usual way

$$\delta\phi = f^\mu\partial_\mu\phi , \tag{2}$$

where f^μ is a conformal Killing vector (i.e, the generator of the transformation). This transformation rule leaves invariant, up to a boundary term, the free field theory ($\lambda = 0$). To achieve the symmetry transformation in the Liouville model the field ϕ must transform as follows

$$\delta\phi = f^\mu\partial_\mu\phi + \frac{1}{\beta}\partial_\mu f^\mu . \tag{3}$$

The Liouville field ϕ can be converted by a canonical transformation into a free field ψ [1]. The aim of this letter is to prove that a similar construction can be made if the auto-interacting term in (1) is replaced by a mass term $\lambda^2\phi^2$ and the scalar field is minimally coupled to a AdS$_2$ backgroud metric [1] with curvature $R = -\lambda^2$. However in this case the transformation mapping the theory into a (massless) free field theory contains an explicit dependence on time. This implies that the free field dynamics is generated by a Hamiltonian which differs from the initial one by a total derivative term in the extended phase-space. As a consequence of our construction the massive field theory is invariant under conformal transformations when the massive field transforms in an involved way which is implicitly defined by the equations of the canonical transformation.

Let us consider the action of a 2-dimensional massive scalar field ϕ minimally coupled to a background metric $g_{\mu\nu}$

$$S = -\frac{1}{2}\int d^2x\sqrt{-g}\left[(\nabla\phi)^2 + \lambda^2\phi^2\right] . \tag{4}$$

If we express the metric in the conformal gauge ($ds^2 = -e^{2\rho}dx^+dx^-$), then we can write

$$\begin{aligned} S &= \int dx^+dx^- \left[\partial_+\phi\partial_-\phi - \frac{\lambda^2}{4}e^{2\rho}\phi^2\right] \\ &= \frac{1}{2}\int dtdx\left[\dot\phi^2 - \phi'^2 - \lambda^2\phi^2e^{2\rho}\right] . \end{aligned} \tag{5}$$

We denote with dot derivation with respect to the time coordinate $t = \frac{1}{2}(x^+ + x^-)$ and with prime derivation with respect to the spatial coordinate $x = \frac{1}{2}(x^+ - x^-)$. The hamiltonian form of the action becomes

---

[1] Actually we will work on the universal covering space CAdS$_2$ to avoid the problem of the existence of closed time-like curves, although we will use the same notation for simplicity

$$S = \int dt dx \left[\dot{\phi} \pi_\phi - \mathcal{H} \right] , \tag{6}$$

where the hamiltonian density $\mathcal{H}$ is

$$\mathcal{H} = \frac{1}{2} \left[\pi_\phi^2 + (\phi')^2 \right] + \frac{\lambda^2}{2} \phi^2 e^{2\rho} . \tag{7}$$

By choosing the background metric to have constant curvature $-\lambda^2$ we obtain

$$R = 8e^{-2\rho} \partial_+ \partial_- \rho = -\lambda^2 , \tag{8}$$

which is the familiar Liouville equation with solution

$$\rho = \frac{1}{2} \log \frac{\partial_+ A_+ \partial_- A_-}{\left(1 + \frac{\lambda^2}{8} A_+ A_- \right)^2} , \tag{9}$$

where $A_\pm(x^\pm)$ are arbitrary chiral functions. We fix the gauge by setting $A_\pm = x^\pm$. The metric with conformal factor (9) is the AdS$_2$ metric with negative curvature $-\lambda^2$ in conformal coordinates $x^\pm$. Now the equation of motion for the field ϕ is

$$\partial_+ \partial_- \phi = -\frac{\lambda^2}{4} e^{2\rho} \phi = -\frac{\lambda^2}{4} \frac{\phi}{\left(1 + \frac{\lambda^2}{8} x^+ x^- \right)^2} . \tag{10}$$

It is easy to check that an arbitrary solution to this equation satisfies

$$\partial_- (e^{2\rho} \partial_+ (e^{-2\rho} \partial_+ \phi)) = 0 , \tag{11}$$

$$\partial_+ (e^{2\rho} \partial_- (e^{2\rho} \partial_- \phi)) = 0 . \tag{12}$$

Let us consider the equation (11). The general solution is

$$\phi = \int^+ \left[e^{2\rho} \int^+ \left(e^{-2\rho} b_+ \right) \right] + a_- \partial_- \rho + b_- , \tag{13}$$

where $a_-(x^-), b_\pm(x^\pm)$ are arbitrary chiral functions. This is not, in general, a solution to the equation (10). However if we set $b_+ = 0$ and insert (13) into (10) we can see that it is a solution provided

$$b_- = \frac{1}{2} \partial_- a_- , \tag{14}$$

and then

$$\phi = \frac{1}{2} \partial_- a_- - \frac{\lambda^2}{8} \frac{x^+ a_-}{1 + \frac{\lambda^2}{8} x^+ x^-} . \tag{15}$$

A similar argument using (12) shows that

$$\phi = -\frac{1}{2} \partial_+ a_+ + \frac{\lambda^2}{8} \frac{x^- a_+}{1 + \frac{\lambda^2}{8} x^+ x^-} , \tag{16}$$

is also a solution to the equation (10). Because the equation is linear the general solution is therefore

$$\phi = -\frac{1}{2}\left(\partial_+ a_+ - \partial_- a_-\right) + \frac{\lambda^2}{8}\frac{a_+ x^- - a_- x^+}{1+\frac{\lambda^2}{8}x^+x^-}\,, \tag{17}$$

with $a_\pm(x^\pm)$ arbitrary chiral functions. It is fairly well-known that the form of the general solution (9) to the Liouville equation induces a canonical transformation between the Liouville theory and a massless free field theory [1]. Mimicking the same procedure we can construct from (17) the following phase-space transformation from the variables ϕ, π_ϕ to the new ones a_+, a_- (which are no longer chiral functions)

$$\phi = -\frac{1}{2}\left(a'_+ + a'_-\right) + \frac{\lambda^2}{8}\frac{a_+ x^- - a_- x^+}{1+\frac{\lambda^2}{8}x^+x^-}\,, \tag{18}$$

$$\begin{aligned}\pi_\phi = &-\frac{1}{2}(a''_+ - a''_-) + \frac{\lambda^2}{8}\frac{a'_+ x^- + a'_- x^+}{1+\frac{\lambda^2}{8}x^+x^-} \\ &+\frac{\lambda^2}{8}\frac{a_+ - a_-}{1+\frac{\lambda^2}{8}x^+x^-} - \frac{\lambda^4}{64}\frac{(x^+ + x^-)(a_+ x^- - a_- x^+)}{(1+\frac{\lambda^2}{8}x^+x^-)^2}\,.\end{aligned} \tag{19}$$

Defining

$$\psi = -\frac{1}{2}(a'_+ + a'_-), \qquad \pi_\psi = -\frac{1}{2}(a''_+ - a''_-)\,, \tag{20}$$

the transformation $\phi, \pi_\phi \to \psi, \pi_\psi$ is canonical up to boundary terms. This can be seen by evaluating the symplectic form

$$\omega = \int dx\ \delta\phi \wedge \delta\pi_\phi\,. \tag{21}$$

Substituting the transformation (18),(19),(20) and after some computation it can be written as (from now on we shall omit the symbol of exterior product)

$$\begin{aligned}\omega = &-\frac{1}{4}\int dx\ \left[\delta a'_+ \delta a''_+ - \delta a'_- \delta a''_-\right] + \int d\left[\frac{1}{4}\delta a'_+ \delta a'_- \right.\\ &\left.+\frac{\lambda^2}{16}\frac{1}{1+\frac{\lambda^2}{8}x^+x^-}\left(x^-\delta a_+ \delta a'_+ - x^-\delta a_+\delta a'_- - x^+\delta a_-\delta a'_+ + x^+\delta a_-\delta a'_-\right)\right] \\ = &\int dx\ \delta\psi\delta\pi_\psi + \omega_b\,,\end{aligned} \tag{22}$$

where ω_b is a boundary term. At this stage we must recall that AdS$_2$ is not a globally hyperbolic space-time, that is, there is no a well-defined Cauchy problem. In order to make the Cauchy problem well-defined we must specify certain boundary conditions of the fields at spatial infinity [2]. These conditions ensure that quantities like the energy of the solutions, the Klein-Gordon product of two solutions to the equation of motion or the symplectic form are conserved (independents of time). Specifically, if we impose that asymptotically $a_\pm$ behaves like

$$a_\pm \sim (x^\pm)^{1-\epsilon} \qquad 0 < \epsilon << 1 \qquad |x^\pm| \to \infty \,, \tag{23}$$

then all the fields fall off to zero rapidly enough to make all the expressions well-defined. With these conditions the bulk part of ω is a convergent expression and on the other hand the boundary contribution ω_b vanishes and then the theory is truly canonically equivalent to the theory of a field ψ which clearly satisfies the (massless) free field equations

$$\dot{\psi} = \pi_\psi, \qquad \dot{\pi}_\psi = \psi'' \,. \tag{24}$$

Because the transformation is time-dependent these equations are not derived from the hamiltonian density (7). To see this explicitly let us consider the following 1-form in the extended phase-space with coordinates $(\phi(x), \pi_\phi(x), t)$

$$\Omega = \int dx \, [\pi_\phi(x)\delta\phi(x)] - H dt \,. \tag{25}$$

Under a canonical transformation to the variables $(\psi(x), \pi_\psi(x), t)$ Ω transforms in the following way

$$\Omega = \int dx \, [\pi_\psi(x)\delta\psi(x)] - \tilde{H} dt + \delta K \,, \tag{26}$$

for some functionals $\tilde{H}$ and K of the extended phase-space coordinates. Then the hamiltonian equations of motion in terms of ψ, π_ψ are derived from the new hamiltonian $\tilde{H}$

$$\dot{\psi} = \{\psi, \tilde{H}\}_{P.B} = \frac{\delta \tilde{H}}{\delta \pi_\psi} \,, \tag{27}$$

$$\dot{\pi}_\psi = \{\pi_\psi, \tilde{H}\}_{P.B} = -\frac{\delta \tilde{H}}{\delta \psi} \,. \tag{28}$$

In the present case this is satisfied with

$$\tilde{H} = \frac{1}{2} \int dx \left[(\psi')^2 + \pi_\psi^2 \right] \,, \tag{29}$$

and

$$\begin{aligned} K = \int dx \Bigg[& -\frac{\lambda^2}{16} \frac{(x^+ + x^-) a'_+ a'_-}{1 + \frac{\lambda^2}{8} x^+ x^-} - \frac{\lambda^2}{16} \frac{a_+ a'_+ - a_- a'_-}{1 + \frac{\lambda^2}{8} x^+ x^-} \\ & + \frac{\lambda^4}{128} \frac{(x^+ + x^-)(a'_+ + a'_-)(a_+ x^- - a_- x^+)}{\left(1 + \frac{\lambda^2}{8} x^+ x^-\right)^2} \\ & + \frac{\lambda^4}{128} \frac{(a_+ x^- - a_- x^+)(a_+ - a_-)}{\left(1 + \frac{\lambda^2}{8} x^+ x^-\right)^2} \\ & - \frac{\lambda^6}{1024} \frac{(x^+ + x^-)(a_+ x^- - a_- x^+)^2}{\left(1 + \frac{\lambda^2}{8} x^+ x^-\right)^3} \Bigg] \,. \end{aligned} \tag{30}$$

K is the time-dependent generating functional of the transformation given as an implicit functional of the extended phase-space variables and the boundary contributions to $\tilde{H}$ and K vanish by virtue of the boundary conditions (23).

We have yet said that the massless free field action

$$S = -\frac{1}{2}\int d^2x\ \partial_\mu\psi\partial^\mu\psi\ , \tag{31}$$

(and therefore the equation of motion) is invariant, up to a boundary term, under the transformation

$$\delta\psi = f^\mu\partial_\mu\psi\ . \tag{32}$$

if f^μ is a conformal Killing vector. The transformation (18),(19),(20) we have just constructed implies that the action (4) and the equation of motion (10) have an implicit hidden symmetry which depends on the coordinates

$$\delta\phi = f^\mu\partial_\mu\psi + \frac{\lambda^2}{8}\frac{x^-\delta a_+ - x^+\delta a_-}{1+\frac{\lambda^2}{8}x^+x^-}\ , \tag{33}$$

where the variations $\delta a_\pm$ are calculated from (20) with $\pi_\psi = \dot{\psi}$ and it must be understood that all the functions which appear in (33) are implicit functionals of ϕ through (18),(19),(20) with $\pi_\phi = \dot{\phi}$. This rather involved symmetry of the equation of motion (10) is the underlying reason which makes possible to find the exact solution (17).

We can understand better the presence of a conformal symmetry in this theory with the following argument. Let us consider the Jackiw-Teitelboim action [3]

$$S = \int d^2x\sqrt{-g}[R+\lambda^2]\phi\ . \tag{34}$$

In this action both the metric and the dilaton ϕ are dynamical fields, so the theory is generically covariant. In conformal gauge, the unconstrained equations of motion are precisely the Liouville equation (8) and the equation for the massive scalar field (10). Both equations are invariant under conformal transformations when ϕ behaves as a scalar (2) and ρ transforms as in (3). Now if we consider the metric to be a fixed background metric solution to the Liouville equation (8), then the equation (10) still maintains the conformal invariance but now ρ remains fixed and ϕ inherits a peculiar transformation law under conformal transformations which is just the one we have described in this letter.

Finally we would like to point out that when we approach to the boundary of the space-time $|x^+x^-| \to \infty$ the solution (17) behaves like

$$\phi = -\frac{1}{2}(\partial_+a_+ - \partial_-a_-) + \frac{a_+}{x^+} - \frac{a_-}{x^-}\ , \tag{35}$$

that is, like a free field which has a simple relation with ψ. So there is a hidden conformal structure in the theory which becomes clear when we approach to the boundary. It would be interesting to see the possible connection of this with the correspondence between field theories in AdS$_{d+1}$ and d-dimensional conformal field theories on the boundary [4]. These questions will be considered in future research.

ACKNOWLEDGMENTS

I want to acknowledge the Generalitat Valenciana for a FPI fellowship. I want to thank J. Navarro-Salas for many useful comments and suggestions and also to K. Bering for discussions and to R. Jackiw for a critical reading of the manuscript.

REFERENCES

[1] E. Braaten, T. Curtright and C. Thorn, *Phys. Lett.* B 118 (1982) 115.
E. D'Hoker and R. Jackiw, *Phys. Rev.* D 26 (1982) 3517.
J. L. Gervais and A. Neveu, *Nucl. Phys.* B 199 (1982) 59.

[2] S. J. Avis, C. J. Isham and D. Storey, *Phys. Rev.* D 18 (1978) 3565.
P. Breitenlohner and D. Z. Freedman, *Ann. Phys.* 37 (1982) 6060.

[3] R. Jackiw and C. Teitelboim, *in Quantum Theory of Gravity,* ed. S. Christensen (Adam Hilger, 1984).

[4] J. Maldacena, *"The Large N Limit of Superconformal field theories and supergravity"*, hep-th/9711200.
E. Witten, *"Anti de Sitter Space and Holography"*, hep-th/9802150.